

Symmetry reductions in loop quantum gravity

based on classical gauge fixings

Norbert Bodendorfer

University of Warsaw

based on arXiv:1410.5608 and
arXiv:1410.5609 (with J. Lewandowski and J. Świeżewski)

Quantum Gravity in Cracow⁴, 09.05.2015

Unterstützt von / Supported by



Alexander von Humboldt
Stiftung/Foundation

Plan of the talk

- 1 Approaches to symmetry reductions
- 2 General strategy
- 3 Bianchi I: Details on classical derivation
- 4 Bianchi I: Details on quantum theory
- 5 Spherical symmetry (sketch)
- 6 Conclusion

Outline

- 1 Approaches to symmetry reductions
- 2 General strategy
- 3 Bianchi I: Details on classical derivation
- 4 Bianchi I: Details on quantum theory
- 5 Spherical symmetry (sketch)
- 6 Conclusion

Proposals for a symmetry reduced quantum theory in LQG

Proposals for a symmetry reduced quantum theory in LQG

- Mini / midi-superspace quantisation

- ▶ LQC [Bojowald '99-; Ashtekar, Bojowald, Lewandowski '03; ...]
- ▶ Schwarzschild black hole [Kastrup, Thiemann '93; Kuchař '94, Gambini, Pullin '13]
- ▶ Spherical symmetry [Bojowald, Kastrup '99, ..., Bojowald, Swiderski '04, ...]
- ▶ More on spherical symmetry [Alvarez, Capurro, Gambini, Pullin, Olmedo, Rastgoo ...]

Proposals for a symmetry reduced quantum theory in LQG

- Mini / midi-superspace quantisation
 - ▶ LQC [Bojowald '99-; Ashtekar, Bojowald, Lewandowski '03; ...]
 - ▶ Schwarzschild black hole [Kastrup, Thiemann '93; Kuchař '94, Gambini, Pullin '13]
 - ▶ Spherical symmetry [Bojowald, Kastrup '99, ..., Bojowald, Swiderski '04, ...]
 - ▶ More on spherical symmetry [Alvarez, Capurro, Gambini, Pullin, Olmedo, Rastgoo ...]
- Approximately symmetric spin networks
 - ▶ Weave states [Ashtekar, Rovelli, Smolin '92; Bombelli '00]
 - ▶ Spinfoam cosmology [Bianchi, Rovelli, Vidotto '10-; Kieselowski, Lewandowski, Puchta '12]
 - ▶ Canonical Bianchi I, reduced states [Alesci, Cianfrani '12-; Pawłowski '14]

Proposals for a symmetry reduced quantum theory in LQG

- Mini / midi-superspace quantisation
 - ▶ LQC [Bojowald '99-; Ashtekar, Bojowald, Lewandowski '03; ...]
 - ▶ Schwarzschild black hole [Kastrup, Thiemann '93; Kuchař '94, Gambini, Pullin '13]
 - ▶ Spherical symmetry [Bojowald, Kastrup '99, ..., Bojowald, Swiderski '04, ...]
 - ▶ More on spherical symmetry [Alvarez, Capurro, Gambini, Pullin, Olmedo, Rastgoo ...]
- Approximately symmetric spin networks
 - ▶ Weave states [Ashtekar, Rovelli, Smolin '92; Bombelli '00]
 - ▶ Spinfoam cosmology [Bianchi, Rovelli, Vidotto '10-; Kieselowski, Lewandowski, Puchta '12]
 - ▶ Canonical Bianchi I, reduced states [Alesci, Cianfrani '12-; Pawłowski '14]
- Study of symmetric connections
 - ▶ Quantisation \leftrightarrow reduction [Bojowald '04; Engle '05; Hanusch '13]
 - ▶ Embedding of states [Engle '07; Brunneemann, Fleischhack '07; Fleischhack '10]

Proposals for a symmetry reduced quantum theory in LQG

- Mini / midi-superspace quantisation
 - ▶ LQC [Bojowald '99-; Ashtekar, Bojowald, Lewandowski '03; ...]
 - ▶ Schwarzschild black hole [Kastrup, Thiemann '93; Kuchař '94, Gambini, Pullin '13]
 - ▶ Spherical symmetry [Bojowald, Kastrup '99, ..., Bojowald, Swiderski '04, ...]
 - ▶ More on spherical symmetry [Alvarez, Capurro, Gambini, Pullin, Olmedo, Rastgoo ...]
- Approximately symmetric spin networks
 - ▶ Weave states [Ashtekar, Rovelli, Smolin '92; Bombelli '00]
 - ▶ Spinfoam cosmology [Bianchi, Rovelli, Vidotto '10-; Kieselowski, Lewandowski, Puchta '12]
 - ▶ Canonical Bianchi I, reduced states [Alesci, Cianfrani '12-; Pawłowski '14]
- Study of symmetric connections
 - ▶ Quantisation \leftrightarrow reduction [Bojowald '04; Engle '05; Hanusch '13]
 - ▶ Embedding of states [Engle '07; Brunnemann, Fleischhack '07; Fleischhack '10]
- Condensate states
 - ▶ GFT [Gielen, Oriti, Sindoni '13-; Calcagni '14, ...]

Proposals for a symmetry reduced quantum theory in LQG

- Mini / midi-superspace quantisation
 - ▶ LQC [Bojowald '99-; Ashtekar, Bojowald, Lewandowski '03; ...]
 - ▶ Schwarzschild black hole [Kastrup, Thiemann '93; Kuchař '94, Gambini, Pullin '13]
 - ▶ Spherical symmetry [Bojowald, Kastrup '99, ..., Bojowald, Swiderski '04, ...]
 - ▶ More on spherical symmetry [Alvarez, Capurro, Gambini, Pullin, Olmedo, Rastgoo ...]
- Approximately symmetric spin networks
 - ▶ Weave states [Ashtekar, Rovelli, Smolin '92; Bombelli '00]
 - ▶ Spinfoam cosmology [Bianchi, Rovelli, Vidotto '10-; Kieselowski, Lewandowski, Puchta '12]
 - ▶ Canonical Bianchi I, reduced states [Alesci, Cianfrani '12-; Pawłowski '14]
- Study of symmetric connections
 - ▶ Quantisation \leftrightarrow reduction [Bojowald '04; Engle '05; Hanusch '13]
 - ▶ Embedding of states [Engle '07; Brunnemann, Fleischhack '07; Fleischhack '10]
- Condensate states
 - ▶ GFT [Gielen, Oriti, Sindoni '13-; Calcagni '14, ...]
- Group averaging w.r.t. symmetry generator at quantum level
 - ▶ Spherical symmetry [NB, Lewandowski, Świeżewski '14]

Proposals for a symmetry reduced quantum theory in LQG

- Mini / midi-superspace quantisation
 - ▶ LQC [Bojowald '99-; Ashtekar, Bojowald, Lewandowski '03; ...]
 - ▶ Schwarzschild black hole [Kastrup, Thiemann '93; Kuchař '94, Gambini, Pullin '13]
 - ▶ Spherical symmetry [Bojowald, Kastrup '99, ..., Bojowald, Swiderski '04, ...]
 - ▶ More on spherical symmetry [Alvarez, Capurro, Gambini, Pullin, Olmedo, Rastgoo ...]
- Approximately symmetric spin networks
 - ▶ Weave states [Ashtekar, Rovelli, Smolin '92; Bombelli '00]
 - ▶ Spinfoam cosmology [Bianchi, Rovelli, Vidotto '10-; Kieselowski, Lewandowski, Puchta '12]
 - ▶ Canonical Bianchi I, reduced states [Alesci, Cianfrani '12-; Pawłowski '14]
- Study of symmetric connections
 - ▶ Quantisation \leftrightarrow reduction [Bojowald '04; Engle '05; Hanusch '13]
 - ▶ Embedding of states [Engle '07; Brunnemann, Fleischhack '07; Fleischhack '10]
- Condensate states
 - ▶ GFT [Gielen, Oriti, Sindoni '13-; Calcagni '14, ...]
- Group averaging w.r.t. symmetry generator at quantum level
 - ▶ Spherical symmetry [NB, Lewandowski, Świeżewski '14]
- Code symmetry as $f(p, q) = 0$, impose $\widehat{f(p, q)} |\Psi\rangle_{\text{sym}} = 0 \leftarrow$ this talk
 - ▶ Bianchi I models [NB '14]
 - ▶ Spherical symmetry [NB, Lewandowski, Świeżewski '14]

Outline

- 1 Approaches to symmetry reductions
- 2 General strategy**
- 3 Bianchi I: Details on classical derivation
- 4 Bianchi I: Details on quantum theory
- 5 Spherical symmetry (sketch)
- 6 Conclusion

General strategy for the symmetry reduction

1 Suitable classical starting point

- ▶ Gauge fix spatial diffeomorphisms adapted to the symmetry reduction
- ▶ Go to the reduced phase space, i.e. solve constraints or employ Dirac bracket
- ▶ Find new connection variables on Γ_{red} (not Ashtekar-Barbero variables)

General strategy for the symmetry reduction

1 Suitable classical starting point

- ▶ Gauge fix spatial diffeomorphisms adapted to the symmetry reduction
- ▶ Go to the reduced phase space, i.e. solve constraints or employ Dirac bracket
- ▶ Find new connection variables on Γ_{red} (not Ashtekar-Barbero variables)

2 Identification of constraints imposed by symmetry reduction

- ▶ Find phase space functions $f_i(p, q) = 0$ in the symmetric subspace
- ▶ $f_i = 0$ may be a first or second class set of constraints
- ▶ Later, choose first class subset via gauge unfixing (\rightarrow Dirac quantisation)

General strategy for the symmetry reduction

1 Suitable classical starting point

- ▶ Gauge fix spatial diffeomorphisms adapted to the symmetry reduction
- ▶ Go to the reduced phase space, i.e. solve constraints or employ Dirac bracket
- ▶ Find new connection variables on Γ_{red} (not Ashtekar-Barbero variables)

2 Identification of constraints imposed by symmetry reduction

- ▶ Find phase space functions $f_i(p, q) = 0$ in the symmetric subspace
- ▶ $f_i = 0$ may be a first or second class set of constraints
- ▶ Later, choose first class subset via gauge unfixing (\rightarrow Dirac quantisation)

3 Quantise the reduced phase space via LQG techniques

- ▶ At this point, still full GR (if accessible by the gauge fixing)
- ▶ At this point, no (full) spatial diffeomorphism constraint at quantum level

General strategy for the symmetry reduction

1 Suitable classical starting point

- ▶ Gauge fix spatial diffeomorphisms adapted to the symmetry reduction
- ▶ Go to the reduced phase space, i.e. solve constraints or employ Dirac bracket
- ▶ Find new connection variables on Γ_{red} (not Ashtekar-Barbero variables)

2 Identification of constraints imposed by symmetry reduction

- ▶ Find phase space functions $f_i(p, q) = 0$ in the symmetric subspace
- ▶ $f_i = 0$ may be a first or second class set of constraints
- ▶ Later, choose first class subset via gauge unfixing (\rightarrow Dirac quantisation)

3 Quantise the reduced phase space via LQG techniques

- ▶ At this point, still full GR (if accessible by the gauge fixing)
- ▶ At this point, no (full) spatial diffeomorphism constraint at quantum level

4 Impose reduction conditions $f_i = 0$ as operator equations: $\hat{f}_i |\Psi\rangle_{\text{sym}} = 0$

- ▶ Find subspace of quantum reduced states $|\Psi\rangle_{\text{sym}}$
- ▶ Find observables \hat{O}_{sym} w.r.t. reduction constraints: $[\hat{O}_{\text{sym}}, \hat{f}_i] = 0$

General strategy for the symmetry reduction

1 Suitable classical starting point

- ▶ Gauge fix spatial diffeomorphisms adapted to the symmetry reduction
- ▶ Go to the reduced phase space, i.e. solve constraints or employ Dirac bracket
- ▶ Find new connection variables on Γ_{red} (not Ashtekar-Barbero variables)

2 Identification of constraints imposed by symmetry reduction

- ▶ Find phase space functions $f_i(p, q) = 0$ in the symmetric subspace
- ▶ $f_i = 0$ may be a first or second class set of constraints
- ▶ Later, choose first class subset via gauge unfixing (\rightarrow Dirac quantisation)

3 Quantise the reduced phase space via LQG techniques

- ▶ At this point, still full GR (if accessible by the gauge fixing)
- ▶ At this point, no (full) spatial diffeomorphism constraint at quantum level

4 Impose reduction conditions $f_i = 0$ as operator equations: $\hat{f}_i |\Psi\rangle_{\text{sym}} = 0$

- ▶ Find subspace of quantum reduced states $|\Psi\rangle_{\text{sym}}$
- ▶ Find observables \hat{O}_{sym} w.r.t. reduction constraints: $[\hat{O}_{\text{sym}}, \hat{f}_i] = 0$

5 Relate observables \hat{O}_{sym} to mini- / midisuperspace parameters

- ▶ Map observables
- ▶ Study dynamics

Outline

- 1 Approaches to symmetry reductions
- 2 General strategy
- 3 Bianchi I: Details on classical derivation**
- 4 Bianchi I: Details on quantum theory
- 5 Spherical symmetry (sketch)
- 6 Conclusion

Classical preparations I: Phase space

Gauge fixing to obtain suitable coordinates

- 1 Start with ADM phase space $\{q_{ab}(\sigma), P^{cd}(\sigma')\} = \delta^{(3)}(\sigma, \sigma') \delta_{(a}^c \delta_{b)}^d$
- 2 Impose **diagonal metric** gauge $q_{a \neq b} = 0 \Leftrightarrow q = \text{diag}(q_{xx}, q_{yy}, q_{zz})$

Classical preparations I: Phase space

Gauge fixing to obtain suitable coordinates

- 1 Start with ADM phase space $\{q_{ab}(\sigma), P^{cd}(\sigma')\} = \delta^{(3)}(\sigma, \sigma') \delta_{(a}^c \delta_{b)}^d$
- 2 Impose **diagonal metric** gauge $q_{a \neq b} = 0 \Leftrightarrow q = \text{diag}(q_{xx}, q_{yy}, q_{zz})$
- 3 Gauge fixes the spatial diffeomorphism constraint $N^a C_a = -2N^a \nabla_b P^b_a = 0$
(up to reduced spatial diffeomorphisms with shift vector $\vec{N} = (N^x(x), N^y(y), N^z(z))$)

Classical preparations I: Phase space

Gauge fixing to obtain suitable coordinates

- 1 Start with ADM phase space $\{q_{ab}(\sigma), P^{cd}(\sigma')\} = \delta^{(3)}(\sigma, \sigma') \delta_{(a}^c \delta_{b)}^d$
- 2 Impose **diagonal metric** gauge $q_{a \neq b} = 0 \Leftrightarrow q = \text{diag}(q_{xx}, q_{yy}, q_{zz})$
- 3 Gauge fixes the spatial diffeomorphism constraint $N^a C_a = -2N^a \nabla_b P^b{}_a = 0$
(up to reduced spatial diffeomorphisms with shift vector $\vec{N} = (N^x(x), N^y(y), N^z(z))$)
- 4 Coordinatise the reduced phase space via $q_{xx}, q_{yy}, q_{zz}, P^{xx}, P^{yy}, P^{zz}$
- 5 Solve $C_a = 0$ for $P^{a \neq b} \Rightarrow P^{a \neq b}(q_{aa}, P^{bb}) \Rightarrow$ insert in Hamiltonian

Classical preparations I: Phase space

Gauge fixing to obtain suitable coordinates

- 1 Start with ADM phase space $\{q_{ab}(\sigma), P^{cd}(\sigma')\} = \delta^{(3)}(\sigma, \sigma') \delta_{(a}^c \delta_{b)}^d$
- 2 Impose **diagonal metric** gauge $q_{a \neq b} = 0 \Leftrightarrow q = \text{diag}(q_{xx}, q_{yy}, q_{zz})$
- 3 Gauge fixes the spatial diffeomorphism constraint $N^a C_a = -2N^a \nabla_b P^b_a = 0$
(up to reduced spatial diffeomorphisms with shift vector $\vec{N} = (N^x(x), N^y(y), N^z(z))$)
- 4 Coordinatise the reduced phase space via $q_{xx}, q_{yy}, q_{zz}, P^{xx}, P^{yy}, P^{zz}$
- 5 Solve $C_a = 0$ for $P^{a \neq b} \Rightarrow P^{a \neq b}(q_{aa}, P^{bb}) \Rightarrow$ insert in Hamiltonian

Choose connection type variables

- 1 Define $e_a e_a = q_{aa}$, $e_a e^a = 1$, without summation, and $E^a = \sqrt{\det q} e^a$
- 2 Define $K_a = K_{ab} e^b$ with K_{ab} being the extrinsic curvature constructed from P^{ab}
- 3 Compute new Poisson brackets: $\{K_a(\sigma), E^b(\sigma')\} = \delta^{(3)}(\sigma, \sigma') \delta_a^b$

Classical preparations I: Phase space

Gauge fixing to obtain suitable coordinates

- 1 Start with ADM phase space $\{q_{ab}(\sigma), P^{cd}(\sigma')\} = \delta^{(3)}(\sigma, \sigma') \delta_{(a}^c \delta_{b)}^d$
- 2 Impose **diagonal metric** gauge $q_{a \neq b} = 0 \Leftrightarrow q = \text{diag}(q_{xx}, q_{yy}, q_{zz})$
- 3 Gauge fixes the spatial diffeomorphism constraint $N^a C_a = -2N^a \nabla_b P^b_a = 0$
(up to reduced spatial diffeomorphisms with shift vector $\vec{N} = (N^x(x), N^y(y), N^z(z))$)
- 4 Coordinatise the reduced phase space via $q_{xx}, q_{yy}, q_{zz}, P^{xx}, P^{yy}, P^{zz}$
- 5 Solve $C_a = 0$ for $P^{a \neq b} \Rightarrow P^{a \neq b}(q_{aa}, P^{bb}) \Rightarrow$ insert in Hamiltonian

Choose connection type variables

- 1 Define $e_a e_a = q_{aa}$, $e_a e^a = 1$, without summation, and $E^a = \sqrt{\det q} e^a$
- 2 Define $K_a = K_{ab} e^b$ with K_{ab} being the extrinsic curvature constructed from P^{ab}
- 3 Compute new Poisson brackets: $\{K_a(\sigma), E^b(\sigma')\} = \delta^{(3)}(\sigma, \sigma') \delta_a^b$
- 4 K_a, E^b are like Ashtekar-Barbero variables without internal indices
 \Rightarrow Abelian gauge theory (Poisson bracket of Maxwell theory)

Classical preparations I: Phase space

Gauge fixing to obtain suitable coordinates

- 1 Start with ADM phase space $\{q_{ab}(\sigma), P^{cd}(\sigma')\} = \delta^{(3)}(\sigma, \sigma') \delta_{(a}^c \delta_{b)}^d$
- 2 Impose **diagonal metric** gauge $q_{a \neq b} = 0 \Leftrightarrow q = \text{diag}(q_{xx}, q_{yy}, q_{zz})$
- 3 Gauge fixes the spatial diffeomorphism constraint $N^a C_a = -2N^a \nabla_b P^b{}_a = 0$
(up to reduced spatial diffeomorphisms with shift vector $\vec{N} = (N^x(x), N^y(y), N^z(z))$)
- 4 Coordinatise the reduced phase space via $q_{xx}, q_{yy}, q_{zz}, P^{xx}, P^{yy}, P^{zz}$
- 5 Solve $C_a = 0$ for $P^{a \neq b} \Rightarrow P^{a \neq b}(q_{aa}, P^{bb}) \Rightarrow$ insert in Hamiltonian

Choose connection type variables

- 1 Define $e_a e_a = q_{aa}$, $e_a e^a = 1$, without summation, and $E^a = \sqrt{\det q} e^a$
- 2 Define $K_a = K_{ab} e^b$ with K_{ab} being the extrinsic curvature constructed from P^{ab}
- 3 Compute new Poisson brackets: $\{K_a(\sigma), E^b(\sigma')\} = \delta^{(3)}(\sigma, \sigma') \delta_a^b$
- 4 K_a, E^b are like Ashtekar-Barbero variables without internal indices
 \Rightarrow Abelian gauge theory (Poisson bracket of Maxwell theory)

At this stage, only Hamiltonian constraint and reduced spatial diffeomorphisms left.

Classical preparations II: Consequences of symmetry

\mathbb{T}^3 Bianchi I universe : 3 scale factors & 3 momenta: $q_{ab}(\sigma) = \text{diag}(q_{xx}, q_{yy}, q_{zz})$

Classical preparations II: Consequences of symmetry

\mathbb{T}^3 Bianchi I universe : 3 scale factors & 3 momenta: $q_{ab}(\sigma) = \text{diag}(q_{xx}, q_{yy}, q_{zz})$

Constraints compatible with a Bianchi I universe

- 1 q_{ab} and P^{ab} are diagonal in suitable coordinates
 \Rightarrow impose $P^{a \neq b}(q_{aa}, P^{bb}) = P^{a \neq b}(K_a, E^b) = 0$

Classical preparations II: Consequences of symmetry

\mathbb{T}^3 Bianchi I universe : 3 scale factors & 3 momenta: $q_{ab}(\sigma) = \text{diag}(q_{xx}, q_{yy}, q_{zz})$

Constraints compatible with a Bianchi I universe

- 1 q_{ab} and P^{ab} are diagonal in suitable coordinates
 \Rightarrow impose $P^{a \neq b}(q_{aa}, P^{bb}) = P^{a \neq b}(K_a, E^b) = 0$
- 2 E^a and K_a are independent of the spatial coordinate in suitable coordinates
 \Rightarrow impose $\partial_a K_b = 0 = \partial_a e_b$

Classical preparations II: Consequences of symmetry

\mathbb{T}^3 Bianchi I universe : 3 scale factors & 3 momenta: $q_{ab}(\sigma) = \text{diag}(q_{xx}, q_{yy}, q_{zz})$

Constraints compatible with a Bianchi I universe

- 1 q_{ab} and P^{ab} are diagonal in suitable coordinates
 \Rightarrow impose $P^{a \neq b}(q_{aa}, P^{bb}) = P^{a \neq b}(K_a, E^b) = 0$
- 2 E^a and K_a are independent of the spatial coordinate in suitable coordinates
 \Rightarrow impose $\partial_a K_b = 0 = \partial_a e_b$

Choose first class subset to impose as strong operator equations

Classical preparations II: Consequences of symmetry

\mathbb{T}^3 Bianchi I universe : 3 scale factors & 3 momenta: $q_{ab}(\sigma) = \text{diag}(q_{xx}, q_{yy}, q_{zz})$

Constraints compatible with a Bianchi I universe

- 1 q_{ab} and P^{ab} are diagonal in suitable coordinates
 \Rightarrow impose $P^{a \neq b}(q_{aa}, P^{bb}) = P^{a \neq b}(K_a, E^b) = 0$
- 2 E^a and K_a are independent of the spatial coordinate in suitable coordinates
 \Rightarrow impose $\partial_a K_b = 0 = \partial_a e_b$

Choose first class subset to impose as strong operator equations

Without proof here, see paper for details:

Classical preparations II: Consequences of symmetry

\mathbb{T}^3 Bianchi I universe : 3 scale factors & 3 momenta: $q_{ab}(\sigma) = \text{diag}(q_{xx}, q_{yy}, q_{zz})$

Constraints compatible with a Bianchi I universe

- 1 q_{ab} and P^{ab} are diagonal in suitable coordinates
 \Rightarrow impose $P^{a \neq b}(q_{aa}, P^{bb}) = P^{a \neq b}(K_a, E^b) = 0$
- 2 E^a and K_a are independent of the spatial coordinate in suitable coordinates
 \Rightarrow impose $\partial_a K_b = 0 = \partial_a e_b$

Choose first class subset to impose as strong operator equations

Without proof here, see paper for details: A maximal first class subset is

- 1 All spatial diffeomorphisms: $\tilde{C}_a[N^a] = \int_{\Sigma} d^3\sigma E^a \mathcal{L}_{\vec{N}} K_a = 0$ (incorporates also reduced ones)
- 2 Abelian Gauß law: $G[\omega] = \int_{\Sigma} d^3\sigma \omega \partial_a E^a = 0$

Classical preparations II: Consequences of symmetry

\mathbb{T}^3 Bianchi I universe : 3 scale factors & 3 momenta: $q_{ab}(\sigma) = \text{diag}(q_{xx}, q_{yy}, q_{zz})$

Constraints compatible with a Bianchi I universe

- 1 q_{ab} and P^{ab} are diagonal in suitable coordinates
 \Rightarrow impose $P^{a \neq b}(q_{aa}, P^{bb}) = P^{a \neq b}(K_a, E^b) = 0$
- 2 E^a and K_a are independent of the spatial coordinate in suitable coordinates
 \Rightarrow impose $\partial_a K_b = 0 = \partial_a e_b$

Choose first class subset to impose as strong operator equations

Without proof here, see paper for details: A maximal first class subset is

- 1 All spatial diffeomorphisms: $\tilde{C}_a[N^a] = \int_{\Sigma} d^3\sigma E^a \mathcal{L}_{\vec{N}} K_a = 0$ (incorporates also reduced ones)
- 2 Abelian Gauß law: $G[\omega] = \int_{\Sigma} d^3\sigma \omega \partial_a E^a = 0$

Result:

Direct consequences of a Bianchi I reduction can be imposed as spatial diffeomorphisms and a Gauß law on the (quantised) reduced phase space (as operator equations).

Classical preparations III: Summary

Phase space: (full GR admitting diagonal metric gauge)

- 1 $K_a(\sigma), E^b(\sigma)$ are 3 + 3 canonical variables per spatial point σ
- 2 Remaining constraints are
 - 1 reduced spatial diffeomorphisms (preserving the diagonal gauge)
 - 2 Hamiltonian constraint

Direct consequences of a reduction to Bianchi I are

- 1 All spatial diffeomorphisms: $\tilde{C}_a[N^a] = \int_{\Sigma} d^3\sigma E^a \mathcal{L}_{\vec{N}} K_a = 0$
- 2 Abelian Gauß law: $G[\omega] = \int_{\Sigma} d^3\sigma \omega \partial_a E^a = 0$

Strategy:

- 1 Quantise full phase space via LQG techniques
- 2 Impose symmetry reduction by imposing $\tilde{C}_a = 0 = G$ at the quantum level

Outline

- 1 Approaches to symmetry reductions
- 2 General strategy
- 3 Bianchi I: Details on classical derivation
- 4 Bianchi I: Details on quantum theory**
- 5 Spherical symmetry (sketch)
- 6 Conclusion

Quantisation I: Full theory in diagonal gauge

Standard LQG type quantisation

- 1 Compute holonomies $h_\gamma^\lambda(K) := \exp\left(i\lambda \int_\gamma K_a ds^a\right)$ and fluxes $E(S) = \int_S E^a d^2s_a$
 γ path, S surface, $\lambda \in \mathbb{Z}$ for $U(1)$, or $\lambda \in \mathbb{R}$ for \mathbb{R}_{Bohr} see e.g. [Corichi, Krasnov '97] for $U(1)$

Quantisation I: Full theory in diagonal gauge

Standard LQG type quantisation

- 1 Compute holonomies $h_\gamma^\lambda(K) := \exp\left(i\lambda \int_\gamma K_a ds^a\right)$ and fluxes $E(S) = \int_S E^a d^2s_a$
 γ path, S surface, $\lambda \in \mathbb{Z}$ for $U(1)$, or $\lambda \in \mathbb{R}$ for \mathbb{R}_{Bohr} see e.g. [Corichi, Krasnov '97] for $U(1)$
- 2 Define positive linear Ashtekar-Lewandowski functional on holonomy-flux algebra
- 3 Representation follows from the GNS construction: Hilbertspace = $L^2(\bar{\mathcal{A}}, d\mu_{\text{AL}})$
 $\bar{\mathcal{A}}$ = generalised $U(1)$ or \mathbb{R}_{Bohr} connections

Quantisation I: Full theory in diagonal gauge

Standard LQG type quantisation

- 1 Compute holonomies $h_\gamma^\lambda(K) := \exp\left(i\lambda \int_\gamma K_a ds^a\right)$ and fluxes $E(S) = \int_S E^a d^2s_a$
 γ path, S surface, $\lambda \in \mathbb{Z}$ for $U(1)$, or $\lambda \in \mathbb{R}$ for \mathbb{R}_{Bohr} see e.g. [Corichi, Krasnov '97] for $U(1)$
- 2 Define positive linear Ashtekar-Lewandowski functional on holonomy-flux algebra
- 3 Representation follows from the GNS construction: Hilbertspace = $L^2(\bar{\mathcal{A}}, d\mu_{\text{AL}})$
 $\bar{\mathcal{A}}$ = generalised $U(1)$ or \mathbb{R}_{Bohr} connections

Remarks

- For \mathbb{R}_{Bohr} : $\lim_{R \rightarrow \infty} \frac{1}{2R} \int_{-R}^R dx f(x) = \int_{\mathbb{R}_{\text{Bohr}}} d\mu_{\text{H}} f(x)$ provides **normalised** and translation invariant Haar measure \Rightarrow per edge: $\mathcal{H} = L^2(\mathbb{R}_{\text{Bohr}}, d\mu_{\text{H}})$

Quantisation I: Full theory in diagonal gauge

Standard LQG type quantisation

- 1 Compute holonomies $h_\gamma^\lambda(K) := \exp\left(i\lambda \int_\gamma K_a ds^a\right)$ and fluxes $E(S) = \int_S E^a d^2s_a$
 γ path, S surface, $\lambda \in \mathbb{Z}$ for $U(1)$, or $\lambda \in \mathbb{R}$ for \mathbb{R}_{Bohr} see e.g. [Corichi, Krasnov '97] for $U(1)$
- 2 Define positive linear Ashtekar-Lewandowski functional on holonomy-flux algebra
- 3 Representation follows from the GNS construction: Hilbertspace = $L^2(\bar{\mathcal{A}}, d\mu_{\text{AL}})$
 $\bar{\mathcal{A}}$ = generalised $U(1)$ or \mathbb{R}_{Bohr} connections

Remarks


- For \mathbb{R}_{Bohr} : $\lim_{R \rightarrow \infty} \frac{1}{2R} \int_{-R}^R dx f(x) = \int_{\mathbb{R}_{\text{Bohr}}} d\mu_{\text{H}} f(x)$ provides **normalised** and translation invariant Haar measure \Rightarrow per edge: $\mathcal{H} = L^2(\mathbb{R}_{\text{Bohr}}, d\mu_{\text{H}})$
- Choosing $\lambda \in \mathbb{Z}$ over $\lambda \in \mathbb{R}$ (i.e. compactifying $\int_\gamma K_a ds^a$) has no justification at this stage (also not later)

Quantisation II: Area operator

Area operator for Abelian theory

- $A(S) = |E(S)| = \left| \int_S E^a d^2 s_a \right|$ is analogous to (absolute value of) electric flux
- Important difference to non-Abelian, e.g. $SU(2)$, area op. $\int_S \sqrt{|E^i E_i|}$:
 - ▶ - Absolute value is **outside** of the integral
 - ▶ - $E(S)$ does not detect closed contractible loops for closed S

While one can also define “non-Abelian like” area operator here, the Abelian one will turn out to be most useful.





Quantisation II: Area operator

Area operator for Abelian theory

- $A(S) = |E(S)| = \left| \int_S E^a d^2 s_a \right|$ is analogous to (absolute value of) electric flux
- Important difference to non-Abelian, e.g. $SU(2)$, area op. $\int_S \sqrt{|E^i E_i|}$:
 - ▶ - Absolute value is **outside** of the integral
 - ▶ - $E(S)$ does not detect closed contractible loops for closed S

While one can also define “non-Abelian like” area operator here, the Abelian one will turn out to be most useful.

- ▶ - Non-trivial topology: $A(S)$ can detect Wilson loops even for closed S ()



$\hat{A}(S)$ measures intersection number $N_{\text{int}} \times \text{rep. label}$: $\hat{A}(S) |h_\gamma^\lambda\rangle = |N_{\text{int}} \lambda| |h_\gamma^\lambda\rangle$

Quantisation III: Imposing the symmetry reduction

Reduction constraints are very familiar from full theory

① All spatial diffeomorphisms: $\tilde{C}_a[N^a] = \int_{\Sigma} d^3\sigma E^a \mathcal{L}_{\vec{N}} K_a = 0$

② Abelian Gauß law: $G[\omega] = \int_{\Sigma} d^3\sigma \omega \partial_a E^a = 0$

⇒ **spatially diffeomorphism invariant** and **gauge invariant** charge (spin) networks!

Quantisation III: Imposing the symmetry reduction

Reduction constraints are very familiar from full theory

- 1 All spatial diffeomorphisms: $\tilde{C}_a[N^a] = \int_{\Sigma} d^3\sigma E^a \mathcal{L}_{\vec{N}} K_a = 0$
- 2 Abelian Gauß law: $G[\omega] = \int_{\Sigma} d^3\sigma \omega \partial_a E^a = 0$

⇒ **spatially diffeomorphism invariant** and **gauge invariant** charge (spin) networks!


Simplest choice of quantum state

Consider spin network made from 3 Wilson loops wrapping around $\mathbb{T}_x^1, \mathbb{T}_y^1, \mathbb{T}_z^1$, meeting in a single vertex v . [c.f. Husain '91, '05]

Mapping to Bianchi I LQC states of

[Ashtekar, Wilson-Ewing '09]

$$|\lambda_x, \lambda_y, \lambda_z\rangle \mapsto |p_1, p_2, p_3\rangle$$



Quantisation III: Imposing the symmetry reduction

Reduction constraints are very familiar from full theory

① All spatial diffeomorphisms: $\tilde{C}_a[N^a] = \int_{\Sigma} d^3\sigma E^a \mathcal{L}_{\vec{N}} K_a = 0$

② Abelian Gauß law: $G[\omega] = \int_{\Sigma} d^3\sigma \omega \partial_a E^a = 0$

⇒ **spatially diffeomorphism invariant** and **gauge invariant** charge (spin) networks!


Simplest choice of quantum state

Consider spin network made from 3 Wilson loops wrapping around $\mathbb{T}_x^1, \mathbb{T}_y^1, \mathbb{T}_z^1$, meeting in a single vertex v . [c.f. Husain '91, '05]

Mapping to Bianchi I LQC states of

[Ashtekar, Wilson-Ewing '09]

$$|\lambda_x, \lambda_y, \lambda_z\rangle \mapsto |p_1, p_2, p_3\rangle$$



Observables w.r.t. the reduction constraints

① Area of closed surfaces → 3 non-trivial areas $A(\mathbb{T}_x^2), A(\mathbb{T}_y^2), A(\mathbb{T}_z^2)$

② Diff-equiv. classes of Wilson loops → 3 non-trivial closed loops along $\mathbb{T}_x^1, \mathbb{T}_y^1, \mathbb{T}_z^1$

Quantisation IV: Dynamics

Hamiltonian constraint / true Hamiltonian (via deparametrisation)

Take original Hamiltonian:

- Evaluate at $q_{a \neq b} = 0$ because of gauge fixing
- Discard $P^{a \neq b}$, $\partial_a e_b$, and $\partial_a K_b$ terms because of reduction constraints

Quantisation IV: Dynamics

Hamiltonian constraint / true Hamiltonian (via deparametrisation)

Take original Hamiltonian:

- Evaluate at $q_{a \neq b} = 0$ because of gauge fixing
- Discard $P^{a \neq b}$, $\partial_a e_b$, and $\partial_a K_b$ terms because of reduction constraints

$\Rightarrow H[N] = \int d^3\sigma N (e_x K_y K_z + e_y K_z K_x + e_z K_x K_y)$ looks like in Bianchi I cosmology

Quantisation IV: Dynamics

Hamiltonian constraint / true Hamiltonian (via deparametrisation)

Take original Hamiltonian:

- Evaluate at $q_{a \neq b} = 0$ because of gauge fixing
- Discard $P^{a \neq b}$, $\partial_a e_b$, and $\partial_a K_b$ terms because of reduction constraints

$\Rightarrow H[N] = \int d^3\sigma N (e_x K_y K_z + e_y K_z K_x + e_z K_x K_y)$ looks like in Bianchi I cosmology

Regularise constraint operator (graph preserving)

- Substitute e_a either by fluxes or Thiemann's trick $e_a = 2\{K_a, V\}$
- Approximate K_a via holonomies: $\int K_a ds^a \approx \sin(\lambda \int K_a ds^a) / \lambda$

Quantisation IV: Dynamics

Hamiltonian constraint / true Hamiltonian (via deparametrisation)

Take original Hamiltonian:

- Evaluate at $q_{a \neq b} = 0$ because of gauge fixing
- Discard $P^{a \neq b}$, $\partial_a e_b$, and $\partial_a K_b$ terms because of reduction constraints

$\Rightarrow H[N] = \int d^3\sigma N (e_x K_y K_z + e_y K_z K_x + e_z K_x K_y)$ looks like in Bianchi I cosmology

Regularise constraint operator (graph preserving)

- Substitute e_a either by fluxes or Thiemann's trick $e_a = 2\{K_a, V\}$
- Approximate K_a via holonomies: $\int K_a ds^a \approx \sin(\lambda \int K_a ds^a) / \lambda$

Choice of λ is crucial! $U(1)$ vs. \mathbb{R}_{Bohr}

- $U(1)$ choice: $\lambda = 1$ gives best approximation
 \Rightarrow "old" LQC dynamics [Ashtekar, Bojowald, Lewandowski '03]

Quantisation IV: Dynamics

Hamiltonian constraint / true Hamiltonian (via deparametrisation)

Take original Hamiltonian:

- Evaluate at $q_{a \neq b} = 0$ because of gauge fixing
- Discard $P^{a \neq b}$, $\partial_a e_b$, and $\partial_a K_b$ terms because of reduction constraints

$\Rightarrow H[N] = \int d^3\sigma N (e_x K_y K_z + e_y K_z K_x + e_z K_x K_y)$ looks like in Bianchi I cosmology

Regularise constraint operator (graph preserving)

- Substitute e_a either by fluxes or Thiemann's trick $e_a = 2\{K_a, V\}$
- Approximate K_a via holonomies: $\int K_a ds^a \approx \sin(\lambda \int K_a ds^a) / \lambda$

Choice of λ is crucial! $U(1)$ vs. \mathbb{R}_{Bohr}

- $U(1)$ choice: $\lambda = 1$ gives best approximation
 \Rightarrow "old" LQC dynamics [Ashtekar, Bojowald, Lewandowski '03]
- \mathbb{R}_{Bohr} allows arbitrarily small $\lambda \in \mathbb{R}$ for better approximation.
"improved" LQC choice: $1/\lambda_x = \sqrt{|E^y E^z / E^x|}$ = size of universe in x -direction
 \Rightarrow "new" LQC dynamics [Ashtekar, Pawłowski, Singh '06; Ashtekar, Wilson-Ewing '09]

Outline

- 1 Approaches to symmetry reductions
- 2 General strategy
- 3 Bianchi I: Details on classical derivation
- 4 Bianchi I: Details on quantum theory
- 5 Spherical symmetry (sketch)**
- 6 Conclusion

Classical preparations for reduction to spherical symmetry

ADM phase space in radial gauge (without details here)


- 1 Start with ADM phase space

Classical preparations for reduction to spherical symmetry

ADM phase space in radial gauge (without details here)

- 1 Start with ADM phase space
- 2 Define adapted coordinate system

[Duch, Kamiński, Lewandowski, Świeżewski '14]



$$(r, \theta, \phi) \leftrightarrow \sigma = \exp_{\sigma_0}(x^l e_l)$$

e_l = specific frame at σ_0 , $l = 1, 2, 3$

x^l = local "cartesian" coordinates


$x^l \leftrightarrow r, \theta, \phi$ spherical coordinates

S_r^2 = "spheres" of constant geodesic distance r

Classical preparations for reduction to spherical symmetry

ADM phase space in radial gauge (without details here)

- 1 Start with ADM phase space
- 2 Define adapted coordinate system
[Duch, Kamiński, Lewandowski, Świeżewski '14]
- 3 Impose **radial** gauge $q_{ra} = \delta_{ra}$



$$(r, \theta, \phi) \leftrightarrow \sigma = \exp_{\sigma_0}(x^l e_l)$$

e_l = specific frame at σ_0 , $l = 1, 2, 3$

x^l = local "cartesian" coordinates

$x^l \leftrightarrow r, \theta, \phi$ spherical coordinates

S_r^2 = "spheres" of constant geodesic distance r

Classical preparations for reduction to spherical symmetry


ADM phase space in radial gauge (without details here)

- 1 Start with ADM phase space
- 2 Define adapted coordinate system
[Duch, Kamiński, Lewandowski, Świeżewski '14]
- 3 Impose **radial** gauge $q_{ra} = \delta_{ra}$
- 4 Express q_{AB}, P^{AB} via

SU(2) connection variables A_A^i, E_j^B

$$A, B = \theta, \phi, \quad \{A_A^i(\sigma), E_j^B(\sigma')\} = \delta^{(3)}(\sigma, \sigma') \delta_A^B \delta_j^i$$

A_A^i, E_j^B = variables of 3d gravity, with spatial slice S_r^2



$$(r, \theta, \phi) \leftrightarrow \sigma = \exp_{\sigma_0}(x^I e_I)$$

e_I = specific frame at σ_0 , $I = 1, 2, 3$

x^I = local "cartesian" coordinates

$x^I \leftrightarrow r, \theta, \phi$ spherical coordinates

S_r^2 = "spheres" of constant geodesic distance r

Classical preparations for reduction to spherical symmetry

ADM phase space in radial gauge (without details here)


- 1 Start with ADM phase space
- 2 Define adapted coordinate system
[Duch, Kamiński, Lewandowski, Świeżewski '14]

- 3 Impose **radial** gauge $q_{ra} = \delta_{ra}$
- 4 Express q_{AB}, P^{AB} via

SU(2) connection variables A_A^i, E_j^B

$A, B = \theta, \phi, \{A_A^i(\sigma), E_j^B(\sigma')\} = \delta^{(3)}(\sigma, \sigma') \delta_A^B \delta_j^i$
 $A_A^i, E_j^B =$ variables of 3d gravity, with spatial slice S_r^2

- 5 Solve $C_a = 0$ for $P^{rr}, P^{rA} \Rightarrow P^{ra}(A_A^i, E_j^B)$
 \Rightarrow insert in Hamiltonian



$$(r, \theta, \phi) \leftrightarrow \sigma = \exp_{\sigma_0}(x^I e_I)$$

$e_I =$ specific frame at $\sigma_0, I = 1, 2, 3$

$x^I =$ local "cartesian" coordinates

$x^I \leftrightarrow r, \theta, \phi$ spherical coordinates

$S_r^2 =$ "spheres" of constant geodesic distance r

Classical preparations for reduction to spherical symmetry


ADM phase space in radial gauge (without details here)

- 1 Start with ADM phase space
- 2 Define adapted coordinate system
[Duch, Kamiński, Lewandowski, Świeżewski '14]
- 3 Impose **radial** gauge $q_{ra} = \delta_{ra}$
- 4 Express q_{AB}, P^{AB} via

SU(2) connection variables A_A^i, E_j^B

$A, B = \theta, \phi, \quad \{A_A^i(\sigma), E_j^B(\sigma')\} = \delta^{(3)}(\sigma, \sigma') \delta_A^B \delta_j^i$
 A_A^i, E_j^B = variables of 3d gravity, with spatial slice S_r^2

- 5 Solve $C_a = 0$ for $P^{rr}, P^{rA} \Rightarrow P^{ra}(A_A^i, E_j^B)$
 \Rightarrow insert in Hamiltonian



$$(r, \theta, \phi) \leftrightarrow \sigma = \exp_{\sigma_0}(x^l e_l)$$

e_l = specific frame at $\sigma_0, \quad l = 1, 2, 3$

x^l = local "cartesian" coordinates

$x^l \leftrightarrow r, \theta, \phi$ spherical coordinates

S_r^2 = "spheres" of constant geodesic distance r

Find conditions compatible with spherical symmetry

- $P^{rA} = 0 \Leftrightarrow$ generator of all spatial diffeomorphisms preserving all S_r^2
 (Follows from non-existence of non-zero spherically symmetric vector field on S^2)

Impose spherical symmetry as invariance under S_r^2 -preserving diffeomorphisms.

(These are **active** diffeomorphisms with respect to the (r, θ, ϕ) coordinate system)

Quantisation and reduction to spherical symmetry

Perform standard LQG-type quantisation (roughly similar to lattice field theory)

- 1 $SU(2)$ gauge theory with holonomies restricted to lie in an S_r^2
 \Rightarrow Quantum states (generalised Wilson loops) have support on different S_r^2

Quantisation and reduction to spherical symmetry

Perform standard LQG-type quantisation (roughly similar to lattice field theory)

- 1 $SU(2)$ gauge theory with holonomies restricted to lie in an S_r^2
 \Rightarrow Quantum states (generalised Wilson loops) have support on different S_r^2
- 2 Reduction constraints lead to spatially diffeomorphism invariance on each S_r^2
(Quantum states "diffeomorphism averaged" over the S_r^2)

Quantisation and reduction to spherical symmetry

Perform standard LQG-type quantisation (roughly similar to lattice field theory)

- 1 $SU(2)$ gauge theory with holonomies restricted to lie in an S_r^2
 \Rightarrow Quantum states (generalised Wilson loops) have support on different S_r^2
- 2 Reduction constraints lead to spatially diffeomorphism invariance on each S_r^2
(Quantum states “diffeomorphism averaged” over the S_r^2)

Observables w.r.t. the reduction constraints

- 1 Areas of the S_r^2 $\rightarrow 4\pi R(r)^2 := \int_{S_r^2} d^2x \sqrt{\det q_{AB}}$
- 2 Averaged trace of momenta $\rightarrow P_R(r) := \frac{2}{R(r)} \int_{S_r^2} d^2x P^{AB} q_{AB}$

(+ all other S_r^2 -preserving diffeomorphism invariant observables. \Rightarrow more than in classically reduced theory!)

Quantisation and reduction to spherical symmetry

Perform standard LQG-type quantisation (roughly similar to lattice field theory)

- 1 $SU(2)$ gauge theory with holonomies restricted to lie in an S_r^2
 \Rightarrow Quantum states (generalised Wilson loops) have support on different S_r^2
- 2 Reduction constraints lead to spatially diffeomorphism invariance on each S_r^2
(Quantum states "diffeomorphism averaged" over the S_r^2)

Observables w.r.t. the reduction constraints

- 1 Areas of the S_r^2 $\rightarrow 4\pi R(r)^2 := \int_{S_r^2} d^2x \sqrt{\det q_{AB}}$
- 2 Averaged trace of momenta $\rightarrow P_R(r) := \frac{2}{R(r)} \int_{S_r^2} d^2x P^{AB} q_{AB}$

(+ all other S_r^2 -preserving diffeomorphism invariant observables. \Rightarrow more than in classically reduced theory!)

What about dynamics?

More challenging than for Bianchi I, ongoing work with A. Zipfel. First steps

- Map states in classically reduced \rightarrow quantum reduced theory
- Compute quantum algebra $[\hat{R}(r), \hat{P}_R(r')]$ from full theory

Outline

- 1 Approaches to symmetry reductions
- 2 General strategy
- 3 Bianchi I: Details on classical derivation
- 4 Bianchi I: Details on quantum theory
- 5 Spherical symmetry (sketch)
- 6 Conclusion**

Conclusion: Proposed reduction programme successful

- 1 **Suitable classical starting point**
 - ▶ Bianchi I: ADM in diagonal metric gauge
 - ▶ Sph. sym.: ADM in radial gauge
- 2 **Identification of constraints** imposed by symmetry reduction
 - ▶ Bianchi I: all spatial diffeomorphisms and Abelian Gauß law
 - ▶ Sph. sym.: S_r^2 -preserving diffeomorphisms (Gauß law from variable choice)
- 3 **Quantise** the reduced phase space via LQG techniques
 - ▶ Bianchi I: Abelian gauge theory
 - ▶ Sph. sym.: SU(2) gauge theory with Wilson loops (graphs) in S_r^2
- 4 **Impose reduction conditions** as operator equations
 - ▶ Bianchi I: spatial diffeomorphism invariance and gauge invariance
 - ▶ Sph. sym.: S_r^2 -preserving spatial diffeomorphism invariance (and gauge invariance)
- 5 **Relate observables** w.r.t. reduction constraints to mini- / midisuperspace
 - ▶ Bianchi I: three areas and conjugate momenta
 - ▶ Sph. sym.: $R(r)$ and $P_R(r)$
- 6 Future work: perturbations to Bianchi I, coarse graining, spherical collapse...

Conclusion: Proposed reduction programme successful

1 Suitable classical starting point

- ▶ Bianchi I: ADM in diagonal metric gauge
- ▶ Sph. sym.: ADM in radial gauge

2 Identification of constraints imposed by symmetry reduction

- ▶ Bianchi I: all spatial diffeomorphisms and Abelian Gauß law
- ▶ Sph. sym.: S_r^2 -preserving diffeomorphisms (Gauß law from variable choice)

3 Quantise the reduced phase space via LQG techniques

- ▶ Bianchi I: Abelian gauge theory
- ▶ Sph. sym.: SU(2) gauge theory with Wilson loops (graphs) in S_r^2

4 Impose reduction conditions as operator equations

- ▶ Bianchi I: spatial diffeomorphism invariance and gauge invariance
- ▶ Sph. sym.: S_r^2 -preserving spatial diffeomorphism invariance (and gauge invariance)

5 Relate observables w.r.t. reduction constraints to mini- / midisuperspace

- ▶ Bianchi I: three areas and conjugate momenta
- ▶ Sph. sym.: $R(r)$ and $P_R(r)$

6 Future work: perturbations to Bianchi I, coarse graining, spherical collapse...

Thank you for your attention!